

SYMPLECTIC AUTOMORPHISMS OF T^*S^2

PAUL SEIDEL

ABSTRACT. Let $T \subset T^*S^2$ be the bundle of unit discs, which is a compact symplectic manifold with boundary. Let $\mathcal{S}$ be the group of symplectic automorphisms of T which are trivial near ∂T . In [5] it was proved that $\pi_0(\mathcal{S})$ contains an element of infinite order. This element is given by the ‘generalized Dehn twist’ τ which is the monodromy map of a quadratic singularity [2] (see also [4]). The fact that $[\tau]$ has infinite order is interesting because the class of τ in $\pi_0(\mathcal{D})$, where $\mathcal{D}$ is the corresponding group of diffeomorphisms, has order two. This note contains a short direct proof of the fact that $\mathcal{S}$ is weakly homotopy equivalent to the discrete group $\mathbb{Z}$, with τ as a generator. The proof proceeds by compactifying T to $S^2 \times S^2$. The topology of the symplectic automorphism group of $S^2 \times S^2$ was determined by Gromov [3], and we use a variant of his argument.

Let $T \subset T^*S^2$ be the space of cotangent vectors of length ≤ 1 , and η its standard symplectic structure. Let $\mathcal{D}$ be the group of diffeomorphisms $\phi : T \rightarrow T$ such that $\phi = \text{id}$ in a neighbourhood of ∂T , with the C^∞ -topology, and $\mathcal{S}$ the subgroup of those ϕ which are symplectic (that is, they preserve η).

Theorem 1. $\pi_0(\mathcal{S}) \cong \mathbb{Z}$, and $\pi_i(\mathcal{S}) = 1$ for all $i > 0$.

The components of $\mathcal{S}$ cannot all be distinguished in $\mathcal{D}$:

Corollary 2. *The image of the map $\pi_0(\mathcal{S}) \rightarrow \pi_0(\mathcal{D})$ is isomorphic to $\mathbb{Z}/2$.*

We will prove both results by compactifying (T, η) to $S^2 \times S^2$ with its standard symplectic structure ω (the one for which both spheres have equal areas). More precisely, one can identify (up to rescaling the symplectic forms) the interior of (T, η) with the complement of the diagonal Δ in $(S^2 \times S^2, \omega)$. This has the following consequence: let $\mathcal{D}_2$ be the group of diffeomorphisms ψ of $S^2 \times S^2$ such that $\psi|_{\Delta} = \text{id}$ and which act trivially on the normal bundle $\nu\Delta$. Then $\mathcal{D}_2$ is w.h.e. (weakly homotopy equivalent) to $\mathcal{D}$. Similarly, the subgroup $\mathcal{S}_2 \subset \mathcal{D}_2$ of those ψ which preserve ω is w.h.e. to $\mathcal{S}$.

Consider the larger group $\mathcal{D}_1 \supset \mathcal{D}_2$ of oriented diffeomorphisms of $S^2 \times S^2$ which are equal to the identity on Δ , and the subgroup $\mathcal{S}_1 \subset \mathcal{D}_1$ of those which are symplectic. Let $\mathcal{G}$ be the group of gauge transformations of $\nu\Delta$

Date: 19/3/1998.

Research supported by NSF grant DMS 9304580.

as an oriented vector bundle, and $\mathcal{G}^s \subset \mathcal{G}$ the subgroup of symplectic gauge transformations. These topological groups form a commutative diagram with exact rows:

$$(1) \quad \begin{array}{ccccccc} 1 & \longrightarrow & \mathcal{S}_2 & \longrightarrow & \mathcal{S}_1 & \longrightarrow & \mathcal{G}^s \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & \mathcal{D}_2 & \longrightarrow & \mathcal{D}_1 & \longrightarrow & \mathcal{G} \longrightarrow 1 \end{array}$$

Lemma 3. *Let $e_x : \mathcal{G}^s \longrightarrow SL_2(\mathbb{R})$ be the evaluation at some point $x \in \Delta$. e_x is a w.h.e.*

Proof. Let $\mathcal{G}^u \subset \mathcal{G}^s$ be the subgroup of unitary gauge transformations. $\mathcal{G}^u$ is a deformation retract of $\mathcal{G}^s$, and $\mathcal{G}_0^u = \mathcal{G}^u \cap \ker(e_x)$ is a deformation retract of $\ker(e_x)$. Now $\mathcal{G}_0^u$ is homeomorphic to the space $\text{Map}_b(S^2, S^1)$ of smooth based maps $S^2 \longrightarrow S^1$. Therefore $\pi_i(\mathcal{G}^u) \cong \pi_0(\text{Map}_b(S^{2+i}, S^1)) \cong H^1(S^{2+i}) = 0$ for all $i \geq 0$. This proves that $\mathcal{G}_0^u$, and hence also $\ker(e_x)$, is weakly contractible. $\square$

Lemma 4. *Let $\iota \in \mathcal{S}_1$ be the involution of $S^2 \times S^2$ which interchanges the two factors. Then the discrete subgroup $\{\text{id}, \iota\} \subset \mathcal{S}_1$ is a deformation retract.*

We postpone the proof of this Lemma to the end of the note. Note that the two components of $\mathcal{S}_1$ are distinguished by the action on homology.

Proof of Theorem 1. We use the long exact sequence of homotopy groups induced by the top row of (1). Lemma 3 and Lemma 4 imply that $\pi_i(\mathcal{S}_2) = 1$ for $i > 1$. For $i = 0$ one obtains a short exact sequence

$$(2) \quad 1 \longrightarrow \pi_1(\mathcal{G}^s) \cong \mathbb{Z} \xrightarrow{\partial} \pi_0(\mathcal{S}_2) \longrightarrow \pi_0(\mathcal{S}_1) \cong \mathbb{Z}/2 \longrightarrow 1.$$

Let $\overline{\Delta} = \{(x, -x) : x \in S^2\} \subset S^2 \times S^2$ be the antidiagonal. Its complement carries the Hamiltonian S^1 -action $\rho(t)(x, y) = (R_{x+y}^t(x), R_{x+y}^t(y))$, where $R_\xi^t \in SO(3)$ denotes the rotation with axis $\xi/|\xi|$ and angle t . The moment map of ρ is $\mu(x, y) = -|x + y|$. Choose an $r \in C^\infty(\mathbb{R}, \mathbb{R})$ with $r(t) = -\pi$ for $t \leq 1/2$ and $r(t) = 0$ for $t \geq 1$. Define $\tau \in \mathcal{S}_2$ by

$$\tau(x, y) = \begin{cases} (y, x) & |x + y| \leq \frac{1}{2}, \\ \rho(r(|x + y|))(x, y) & \text{otherwise.} \end{cases}$$

Since $\tau|\overline{\Delta} = \iota|\overline{\Delta}$ reverses the orientation of $\overline{\Delta}$, τ acts nontrivially on homology. Hence $[\tau] \in \pi_0(\mathcal{S}_2)$ maps to the nontrivial element of $\pi_0(\mathcal{S}_1)$. τ^2 can be deformed to the identity map by a path $(h_s)_{0 \leq s \leq 1}$ in $\mathcal{S}_1$, e.g.

$$h_s(x, y) = \begin{cases} (x, y) & |x + y| \leq \frac{1}{2}, \\ \rho(2s(\pi + r(|x + y|))) & \text{otherwise.} \end{cases}$$

The derivative of h_s along Δ is a rotation of $\nu\Delta$ with angle $2\pi s$. It follows that $[\tau]^2 = \partial(r)$ where $r \in \pi_1(\mathcal{G}^s)$ is a generator. In view of (2) this shows that $\pi_0(\mathcal{S}_2) \cong \mathbb{Z}$, with $[\tau]$ as a generator. $\square$

In the terminology of [5] τ is the generalized Dehn twist along the Lagrangian sphere $\overline{\Delta}$. Since we can identify $\text{int}(T)$ with $S^2 \times S^2 \setminus \Delta$ in such a way that $\overline{\Delta}$ corresponds to the zero-section $Z \subset T^*S^2$, it follows that $\pi_0(\mathcal{S})$ is generated by the generalized Dehn twist along Z .

Proof of Corollary 2. Consider the diagram of induced maps

$$\begin{array}{ccccccc} 1 & \longrightarrow & \pi_1(\mathcal{G}^s) \cong \mathbb{Z} & \xrightarrow{\partial} & \pi_0(\mathcal{S}_2) \cong \mathbb{Z} & \longrightarrow & \pi_0(\mathcal{S}_1) \cong \mathbb{Z}/2 \\ & & \alpha_1 \downarrow & & \alpha_2 \downarrow & & \alpha_3 \downarrow \\ \pi_1(\mathcal{D}_1) & \longrightarrow & \pi_1(\mathcal{G}) & \xrightarrow{\partial'} & \pi_0(\mathcal{D}_2) & \longrightarrow & \pi_0(\mathcal{D}_1). \end{array}$$

α_3 is injective, and since $\mathcal{G}^s$ is a deformation retract of $\mathcal{G}$, α_1 is an isomorphism. It remains to show that $\partial' = 0$, or equivalently, to find a loop $(\lambda_t)_{0 \leq t \leq 1}$ in $\mathcal{D}_1$ whose action on $\nu\Delta$ is a generator of $\pi_1(\mathcal{G})$. Such a loop is given by $\lambda_t(x, y) = (R_y^{2\pi t}(x), y)$. $\square$

Proof of Lemma 4. A map $\psi \in \mathcal{D}_1$ can act on $H_2(S^2 \times S^2)$ in two ways: either trivially or, like ι , by interchanging the two generators. Let $\mathcal{D}_+ \subset \mathcal{D}_1$ be the subgroup of those ψ which act trivially, and $\mathcal{S}_+$ the corresponding subgroup of $\mathcal{S}_1$. It is sufficient to show that $\mathcal{S}_+$ is contractible. To do this we use an argument due to Gromov [3, 2.4.A₁] (see [1] for further applications of this kind of argument). Let $\mathcal{J}$ be the space of those ω -compatible almost complex structures on $S^2 \times S^2$ whose restriction to $T\Delta$ agrees with that of the standard complex structure J_0 . There is a map $A : \mathcal{S}_+ \rightarrow \mathcal{J}$ given by $A(\phi) = \phi^*(J_0)$. Gromov's argument provides a map $B : \mathcal{J} \rightarrow \mathcal{S}_+$ such that $B \circ A = \text{id}$; since $\mathcal{J}$ is contractible, this proves that $\mathcal{S}_+$ is also contractible.

Recall that any $J \in \mathcal{J}$ determines, through its pseudo-holomorphic curves, a pair $(\mathcal{F}_1, \mathcal{F}_2)$ of two-dimensional symplectic foliations of $S^2 \times S^2$, which are transverse to each other. The leaves of these foliations are two-spheres whose homology class is $A_1 = [S^2 \times \text{point}]$ resp. $A_2 = [\text{point} \times S^2]$. Since Δ is J -holomorphic and $[\Delta] \cdot A_i = 1$, any leaf of $\mathcal{F}_1$ or $\mathcal{F}_2$ intersects Δ transversely in a single point. It follows that there is a unique $\phi \in \mathcal{D}_+$ which maps $\mathcal{F}_1$ and $\mathcal{F}_2$ to the two standard foliations of $S^2 \times S^2$ by two-spheres. ϕ is not symplectic, but since $\phi^*\omega$ is tamed by J , one can use Moser's technique to deform ϕ into a symplectic automorphism ϕ' ; this can be done in such a way that $\phi' \in \mathcal{S}_+$. Since $\phi = \phi(J)$ depends smoothly on J , one can arrange that $\phi' = \phi'(J)$ also depends smoothly on J , and that $\phi'(J) = \phi(J)$ for all those J for which ϕ is already symplectic. Then one defines B by $B(J) = \phi'(J)$. $\square$

Remark 5. The higher-dimensional analogue of the problem which we have considered presents some potentially more subtle features. Let T_n be the

bundle of unit discs in T^*S^n , and $\mathcal{S}_n$ the group of symplectic automorphisms of T_n which are the identity near the boundary. Using a generalization of the argument in [5] it can be shown that $\pi_0(\mathcal{S}_n)$ contains a class $[\tau]$ of infinite order. However, there is a possible source for elements other than powers of $[\tau]$. The group $\pi_0(\text{Diff}^+(S^n))$, which is isomorphic to the group of exotic $(n+1)$ -spheres for $n \geq 4$, acts on $\pi_0(\mathcal{S}_n)$ by conjugation, and it is not known whether this action preserves $[\tau]$.

REFERENCES

- [1] M. Abreu, *Topology of symplectomorphism groups of $S^2 \times S^2$* , Invent. Math. **131** (1998), 1–24.
- [2] V. I. Arnol'd, *Some remarks on symplectic monodromy of Milnor fibrations*, The Floer Memorial Volume (H. Hofer, C. Taubes, A. Weinstein, and E. Zehnder, eds.), Progress in Mathematics, vol. 133, Birkhäuser, 1995, pp. 99–104.
- [3] M. Gromov, *Pseudoholomorphic curves in symplectic manifolds*, Invent. Math. **82** (1985), 307–347.
- [4] P. Seidel, *Floer homology and the symplectic isotopy problem*, Ph.D. thesis, Oxford University, 1997.
- [5] ———, *Lagrangian two-spheres can be symplectically knotted*, Preprint, 1998.

INSTITUTE FOR ADVANCED STUDY, PRINCETON

E-mail address: pseidel@math.ias.edu